

E1

Electroporation of cells and tissues - interactive e-learning course

Selma Čorović, Samo Mahnič-Kalamiza

University of Ljubljana, Faculty of electrical engineering

Duration of the experiment: app. 90 min

Max. number of participants: 18

Location: Computer room (P9-B0)

Level: Basic

PREREQUISITES

No specific knowledge is required for this laboratory practice.

The aim of this laboratory practice is to provide the participants with basic knowledge on local electric field distribution in cells and tissues exposed to high voltage electric pulses (i.e. electroporation pulses) by means of interactive e-learning course content. The e-learning content is based on the available knowledge from scientific literature.

PROTOCOL OF THE E-LEARNING COURSE

The participants will be gathered in a computer-computer classroom providing each participant with a computer. A short test will be given to establish the baseline knowledge before the e-learning course.

Within the first part of the e-learning course we will bring together the educational material on basic mechanisms underlying electroporation process on the levels of cell membrane, cell and tissues as a composite of cells (Figure 1) [1-2].




Figure 1: Introduction of small molecules through a cell membrane (a) into an electroporated cell (b) and into the successfully electroporated cells within an exposed tissue (c) [9].

Within the second part of the course we will provide basic knowledge on important parameters of local electric field needed for efficient cells and tissue electroporation [3-7], such as: electrode geometry (needle or plate electrodes as illustrated in Figure 2, electrode position with respect to the target tissue and its surrounding the tissues (Figure 3), the contact surface between the electrode and the tissue, the voltage applied to the electrodes and electroporation threshold values. This part of the e-learning course content will be provided by an interactive module we developed to visualize the local electric field distribution in 2D and 3D dimensional tissue models.

The objective of this module is to provide:

- local electric field visualization in cutaneous (protruding tumors) and subcutaneous tumors (tumors more deeply seeded in the tissue);
- guideline on how to overcome a highly resistive skin tissue to permeabilize more conductive underlying tissues and
- visualization and calculation of successfully electroporated volume of the target tissue and its surrounding tissue (i.e. the treated tissue volume exposed to the electric field between reversible and irreversible electroporation threshold value $E_{rev} \leq E < E_{irrev}$) with respect to the selected parameters such as: number and position of electrodes, applied voltage on the electrodes.




Figure 2: Plate electrodes vs. needle electrodes with respect to the target tissue (e.g. tumor tissue).




Figure 3: Electric field distribution within the tumor (inside the circle) and within its surrounding tissue (outside the circle) obtained with three different selection of parameters (number and position of electrodes and voltage applied): (a) 4 electrodes, (b) 8 electrodes and (c) 8 electrodes with increased voltage on electrodes so that the entire volume of tumor is exposed to the $E_{rev} \leq E < E_{irrev}$ [8-10].

After the e-learning course the pedagogical efficiency of presented educational content and the e-learning application usability will be evaluated.

REFERENCES:

- [1] Serša G., Miklavcic D. Electrochemotherapy of tumours (Video Article). *J Visual Exp* 22:1038, 2008.
- [2] Čorović S., Mir L.M., Miklavčič D. In vivo muscle electroporation threshold determination: realistic numerical models and in vivo experiments. *J Memb Biol*, 245:509-520, 2012.
- [3] Čorović S., Županič A., Miklavčič D. Numerical modeling and optimization of electric field distribution in subcutaneous tumor treated with electrochemotherapy using needle electrodes. *IEEE T Plasma Sci*, 36:1665-1672, 2008.

- [4] Čorović S., Županič A., Kranjc S., Al Sakere B., Leroy-Willig A., Mir L.M., Miklavčič D. The influence of skeletal muscle anisotropy on electroporation: in vivo study and numerical modeling. *Med Biol Eng Comput*, 48:637-648, 2010.
- [5] Neal R.E. II, Garcia P.A., Robertson J.L., Davalos R.V. Experimental characterization and numerical modeling of tissue electrical conductivity during pulsed electric fields for irreversible electroporation treatment planning. *IEEE T Biomed Eng*, 59:1077-1085, 2012.
- [6] Čorović S., Pavlin M., Miklavčič D. Analytical and numerical quantification and comparison of the local electric field in the tissue for different electrode configurations. *Biomed Eng Online*, 6:37, 2007.
- [7] Essone Mezeme M., Pucihar G., Pavlin M., Brosseau C., Miklavčič D. A numerical analysis of multicellular environment for modeling tissue electroporation. *Appl Phys Lett*, 100:143701, 2012.
- [8] Čorović S., Lacković I., Šuštarčič P., Šuštar T., Rodič T., Miklavčič D. Modeling of electric field distribution in tissues during electroporation. *Biomed Eng Online*, 12:16, 2013.
- [9] Čorović S., Bešter J., Miklavčič D. An e-learning application on electrochemotherapy. *Biomed Eng Online*, 8:26, 2009.
- [10] Mahnič-Kalamiza S., Kotnik T., Miklavčič D. Educational application for visualization and analysis of electric field strength in multiple electrode electroporation. *BMC Med Educ*, 12:102, 2012.

NOTES & RESULTS
